

# Getting Results with Negative Thinking

D. Loeb

LaBRI, URA CNRS 1304  
Université de Bordeaux I  
33405 Talence, France

E. Damiani and O. D'Antona

Dipartimento di Scienze dell'Informazione  
Università di Milano  
Milano, Italy

February 8, 2008

## Abstract

Given a universe of discourse  $U$ , a *multiset* can be thought of as a function  $M$  from  $U$  to the natural numbers  $\mathbf{N}$ . In this paper, we define a *hybrid set* to be any function from the universe  $U$  to the integers  $\mathbf{Z}$ . These sets are called hybrid since they contain elements with either a positive or negative multiplicity.

Our goal is to use these hybrid sets *as if* they were multisets in order to adequately generalize certain combinatorial facts which are true classically only for nonnegative integers. However, the definition above does not tell us much about these hybrid sets; we must define operations on them which provide us with the needed combinatorial structure.

In section 1, we will define an analog of a set which can contain either a positive or negative number of elements. We will allow sums to be calculated over an arbitrary hybrid set and in particular over “improper” intervals. This will lead us in section 2 to the calculation of the *complementary symmetric functions*  $\text{comp}_n(H)$  whose argument is a hybrid set of variables which is at the same time a generalization of elementary and complete symmetric functions.

In particular, as we will see in section 2.2, the complementary symmetric function generalizes the following two classic results concerning sequences of polynomials with *persistant roots*, that is to say, having the form  $p_n(x) = (x - a_1)(x - a_2) \cdots (x - a_n)$ .

$$p_n(x) = \sum_{k=0}^{\infty} e_k(-a_1, \dots, -a_n) x^{n-k} \quad (1)$$

$$x^n = \sum_{k=0}^{\infty} h_k(a_1, \dots, a_{n-k+1}) p_{n-k}(x). \quad (2)$$

The complementary symmetric function not only allows us to express the connection constants between two arbitrary sequences of polynomials with persistant roots, but to also expand arbitrary rational functions in terms of “extended” sequences with persistant roots and persistant poles.

Many sequences of combinatorial interest  $c_{nk}$  can be defined in terms of connection constants. The complementary symmetric function therefore allows us to simultaneously study them: extending them to the case where  $n$  and  $k$  might be negative, demonstrating their recursions, characterizing them with simple formulas, and providing general inversion formulas. Along these lines, in section 3, we calculate generalized binomial coefficients and give them a combinatorial interpretation in terms of a partial order on the hybrid sets. Similarly, we generalize the Gaussian coefficients, and explain their combinatorial significance. Finally, in section 5, we study a generalization of the  $p, q$ -Stirling numbers of the first and second kind.

# 1 Hybrid Sets

## 1.1 Notation

One usually thinks of a multiset  $M$  as a function from some universe  $U$  to the natural numbers  $\mathbf{N}$ .  $M(x)$  is called the *multiplicity* of  $x$  in the multiset  $M$ . Sets then are a special case of multisets in which all  $x$  have multiplicity either 1 or 0.

In this paper, we generalize these concepts to new “sets” and new “multisets” which may have negative integers as multiplicities as well as nonnegative integers.

**Definition 1 (Hybrid Sets)** *Given a universe  $U$ , any function  $f : U \rightarrow \mathbf{Z}$  is called a hybrid set.*

*The value of  $f(x)$  is said to be the multiplicity of the element  $x$ . If  $f(x) \neq 0$  we say  $x$  is a member of  $f$  and write  $x \in f$ ; otherwise, we write  $x \notin f$ . If the support of  $f$  is finite, define the number of elements  $\#f$  to be the sum  $\sum_{x \in U} f(x)$ .  $f$  is said to be an  $\#f$ -element hybrid set.*

*Given two hybrid sets  $f$  and  $g$ , their sum or union is written as the hybrid set  $f + g$  for which the multiplicity of  $x$  is the sum of the multiplicities in  $f$  and  $g$ :  $(f + g)(x) = f(x) + g(x)$ . Similarly, the hybrid set difference of  $f$  and  $g$  is denoted  $f - g$  and is given by  $(f - g)(x) = f(x) - g(x)$ .*

We will denote hybrid sets by using the usual set braces  $\{\}$  and inserting a bar in the middle  $\{|\}$ . Elements occurring with positive multiplicity are written on the left of the bar, and elements occurring with negative multiplicity are written on the right. Order is completely irrelevant.

For example, if  $f = \{a, b, c, b|d, e, e\}$  then  $f(a) = 1$ ,  $f(b) = 2$ ,  $f(c) = 1$ ,  $f(d) = -1$ ,  $f(e) = -2$ , and  $f(x) = 0$  for  $x \neq a, b, c, d, e$ .

So far, we haven’t accomplished much. These hybrid sets are equivalent to elements of the free group over the universe in question, so one might wonder if there is anything new here. However, we are planning on treating these hybrid sets as if they were multisets. That is to say, we will consider the subsets of hybrid sets; we will perform sums over hybrid sets; and we will enumerate certain classes of combinatorial objects with them. We begin by considering which hybrid sets correspond to the classical notion of a set.

**Definition 2 (New Sets)** *A positive or classical set is a hybrid set taking only values 0 and 1. A negative set is a hybrid set taking only values 0 and  $-1$ . A new set is either a positive or a negative set.*

For example, the set  $S = \{a, b, c\}$  can be written as the classical or positive set  $S = \{b, a, c|\}$  while  $-S = \{|\{a, b, c\}$  is a negative set. Both  $S$  and  $-S$  are new sets. However,  $\{a|b\}$  is not a new set.

The *empty set*  $\emptyset = \{|\}$  is the unique hybrid set for which all elements have multiplicity zero. It is thus simultaneously a positive set and a negative set.

## 1.2 Sums and Products over a Hybrid Set

One often wishes to sum an expression over all the elements of a *set*. For example, we usually write

$$\phi(n) = \sum_{i \in \text{fact}(n)} i$$

for the Euler phi function where  $\text{fact}(n)$  is the set of positive factors of an integer  $n$ .

When dealing with *multisets*, it is often convenient to include multiplicities in the sum or product. One defines  $\sum_{x \in M} F(x)$  and  $\prod_{x \in M} F(x)$  to be  $\sum_{x \in U} M(x)F(x)$  and  $\prod_{x \in U} F(x)^{M(x)}$  respectively. For example, we can expand a monic polynomial  $p(x)$  by

$$p(x) = \prod_{\lambda \in \text{Roots}(p)} (x - \lambda)$$

where  $\text{Roots}(p)$  is the multiset of roots of the polynomial  $p$ .

This notation can be immediately generalized in the obvious way to *hybrid sets*.

**Definition 3 (Sums and Products over Hybrid Sets)** *For any function  $F$  and hybrid set  $h$  we write  $\sum_{x \in h} F(x)$  for  $\sum_{x \in U} h(x)F(x)$ , and we write  $\prod_{x \in h} F(x)$  for  $\prod_{x \in U} F(x)^{h(x)}$ .*

For example, consider the following extension of the definition of  $\text{Roots}(p)$ .

**Definition 4 (Hybrid Set of Roots)** *Let  $f(x)$  be a rational function. Then  $f(x)$  can be written as*

$$f(x) = c \frac{(x - a_1)(x - a_2) \cdots (x - a_i)}{(x - b_1)(x - b_2) \cdots (x - b_j)}.$$

*We then call*

$$\text{Roots}(f) = \{a_1, a_2, \dots, a_i | b_1, b_2, \dots, b_j\}$$

*the Hybrid Set of Roots of  $f(x)$*

For  $f(x)$  a monic rational function, we thus have

$$f(x) = \prod_{\lambda \in \text{Roots}(f)} (x - \lambda).$$

### 1.3 Sums and Products with Limits

Often, sets are written using ellipsis for missing elements. For example,  $\{a_1, \dots, a_5\}$  stands for the set  $\{a_1, a_2, a_3, a_4, a_5\}$ . In the case of hybrid sets, this notation turns out to be rather powerful.

**Definition 5 (Ellipsis)** Let  $(a_n)_{n \in \mathbb{Z}}$  be a sequence of constants indexed by an integer  $n$ . Their ellipsis is defined by the recursion:

$$\begin{aligned} \{a_1..a_0\} &= \emptyset = \{|\} \\ \{a_1..a_n\} &= \{a_1..a_{n-1}\} + \{a_n|\}. \end{aligned}$$

It therefore follows that  $\{a_i..a_j\}$  is explicitly given by:

$$\{a_i..a_j\} = \begin{cases} \{a_i, a_{i+1}, \dots, a_{j-1}, a_j|\} & \text{if } i \leq j, \\ \emptyset & \text{if } i = j + 1, \text{ and} \\ \{|a_{j+1}, a_{j+2}, \dots, a_{i-1}\} & \text{if } i \geq j + 2; \end{cases}$$

In the first case, we are reduced to the standard usage of ellipsis. However, in the other cases, the interval being considered is “improper” classically, but the meaning is clear here. Reference is being made to a certain  $j - i + 1$ -element negative set. Our ellipsis are written with only two dots to distinguish them from normal ellipsis.

One often uses the notation  $\sum_{n=1}^i A(n)$  where  $n$  is a nonnegative integer to denote the sum of  $A(n)$  over the set  $\{1, 2, \dots, n\}$ . Using the ellipsis notation, we can define the sum or product of a quantity from an arbitrary integer to another:

**Definition 6 (Sums and Products with Limits)** Let  $(A(n))_{n \in \mathbb{Z}}$  be a sequence of constants indexed by an integer  $n$ , and let  $i$  and  $j$  be arbitrary integers. The sum and product of  $A(n)$  from  $i$  to  $j$  is given by

$$\begin{aligned} \sum_{n=i}^j A(n) &= \sum_{n \in \{i..j\}} A(n) \\ \prod_{n=i}^j A(n) &= \prod_{n \in \{i..j\}} A(n). \end{aligned}$$

Just as with integrals, the sum is shift invariant, the sum over an empty interval is zero, and the sum over a positive expression over an “improper” interval  $j < i$  is negative.

For example,  $\sum_{i=1}^4 i = 1 + 2 + 3 + 4 = 10$  since  $\{1..4\} = \{1, 2, 3, 4|\}$ , and  $\sum_{i=1}^{-4} = -(-3) - (-2) - (-1) - 0 = 6$  since  $\{1..-4\} = \{|0, -1, -2, -3\}$ . In general,  $\sum_{i=1}^n i = n(n+1)/2$  for all  $n$  regardless of sign. In fact,

**Proposition 7** Let  $p(x)$  be any polynomial, then there is a polynomial  $q(x)$  such that  $q(n) = \sum_{i=1}^n p(i)$  for all  $n$  regardless of sign.  $\square$

*Proof:* Classical proof by induction can be applied here *mutatis mutandis*.  $\square$

Finally, note the following reciprocity formula useful in calculations with improper intervals:

**Proposition 8** Let  $(a_i)_{i \in \mathbb{Z}}$  be a sequence of constants indexed by an integer  $i$ , then for all integers  $n$

$$\{a_1..a_n\} = -\{b_1..b_{-n}\}$$

where  $b_i = a_{n+i}$ .  $\square$

## 2 The Complementary Symmetric Function

### 2.1 Introduction

The complementary symmetric function is so called because it *complements* or extends the *complete* and *elementary* symmetric functions. Recall that  $e_n(V)$  and  $h_n(V)$ , the elementary and complete symmetric functions of degree  $n$  over the set of variables  $V$ , are given by the following generating functions:

$$\begin{aligned}\sum_{n \geq 0} h_n(V) t^n &= \prod_{x \in V} (1 - xt)^{-1} \\ \sum_{n \geq 0} e_n(V) t^n &= \prod_{x \in V} (1 + xt).\end{aligned}$$

The *complementary symmetric function* combines these two definitions by using our notion of the product over a hybrid set.

**Definition 9** *Let  $V$  be a hybrid set of variables. Then the generating function for the complete symmetric function of degree  $n$  over these variables is given by the product:*

$$\sum_{n \geq 0} \text{comp}_n(V) t^n = \prod_{x \in V} (1 - xt).$$

In other words, we have

$$\sum_{n \geq 0} \text{comp}_n(a_1, a_2, \dots | b_1, b_2, \dots) t^n = \frac{\prod_{i=1}^{\infty} (1 - a_i t)}{\prod_{i=1}^{\infty} (1 - b_i t)} = \frac{\sum_{n=0}^{\infty} e_n(-a_1, -a_2, \dots) t^n}{\sum_{n=0}^{\infty} h_n(a_1, a_2, \dots) t^n}.$$

In the case where  $V$  is a negative or positive set, it follows that

$$e_n(a, b, c, \dots) = \text{comp}_n(-a, -b, -c, \dots) \quad (3)$$

$$h_n(a, b, c, \dots) = \text{comp}_n(a, b, c, \dots). \quad (4)$$

It is now immediate that the transformation

$$* : e_n \mapsto h_n$$

used in the theory of symmetric functions is an involution.  $*$  acts on the set of symmetric variables, inverting the sign and the multiplicity of each variable.

The next few results are based on the following convolution identity.

**Lemma 10** *Suppose  $V$  and  $W$  are two (hybrid) sets of variables. Then*

$$\text{comp}_n(V + W) = \sum_{k=0}^n \text{comp}_k(V) \text{comp}_{n-k}(W).$$

*Proof:* Using the definition of  $V + W$  (Definition 1), we can proceed as follows

$$\begin{aligned}
\sum_{n=0}^{\infty} \text{comp}_n(V + W)t^n &= \prod_{x \in V+W} (1 - xt) \\
&= \left( \prod_{x \in V} (1 - xt) \right) \left( \prod_{x \in W} (1 - xt) \right) \\
&= \left( \sum_{n=0}^{\infty} \text{comp}_n(V)t^n \right) \left( \sum_{n=0}^{\infty} \text{comp}_n(W)t^n \right) \\
&= \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \text{comp}_k(V) \text{comp}_{n-k}(W) \right) t^n,
\end{aligned}$$

and identify coefficients of  $t^n$ .  $\square$

We have the following explicit characterization of the complementary symmetric function.

**Theorem 11** *The complementary symmetric function  $\text{comp}_n(a_1, a_2, \dots | b_1, b_2, \dots)$  is the sum of all monomials in the  $a$ 's and the  $b$ 's in which the  $a$ 's may appear with no exponent two or greater, and are associated with a minus sign.*

For example,

$$\begin{aligned}
\text{comp}_2(a, b, c | x, y, z) &= ab + ac + bc \\
&\quad - ax - ay - az - bx - by - bz - cx - cy - cz \\
&\quad + x^2 + xy + xz + y^2 + yz + z^2.
\end{aligned}$$

*Proof of Theorem 11:* First, by Lemma 10,

$$\text{comp}_n(a_1, a_2, \dots | b_1, b_2, \dots) = \sum_{k=0}^n \text{comp}_k(a_1, a_2, \dots) \text{comp}_{n-k}(b_1, b_2, \dots).$$

Then by equations 3 and 4, we have in turn

$$\text{comp}_n(a_1, a_2, \dots | b_1, b_2, \dots) = \sum_{k=0}^n e_k(-a_1, -a_2, \dots) h_{n-k}(b_1, b_2, \dots)$$

where as we recall the elementary symmetric function is the sum of monomials with no repeated variables, and the complete symmetric function is the sum of monomials possibly with repeated variables.  $\square$

The complementary symmetric function satisfies an interesting recursion:

**Theorem 12** *For all  $n, m, k$  positive integers, we have*

$$\begin{aligned}
\text{comp}_n(a_1, \dots, a_m | b_1, \dots, b_k) &= \text{comp}_n(a_1, \dots, a_{m-1} | b_1, \dots, b_{k-1}) \\
&\quad + (b_k - a_m) \text{comp}_{n-1}(a_1, \dots, a_{m-1} | b_1, \dots, b_k). \square
\end{aligned}$$

*Proof:* Either apply Lemma 10 first with  $W = \{a_m|b_k\}$  and then again with  $W = \{a_m|\}$ , or reason with Equations 3 and 4.  $\square$

Note that Theorem 12 holds equally well for  $m, k \leq 0$  provided that  $\text{comp}_n(a_1, \dots, a_m|b_1, \dots, b_k)$  is read as the complementary symmetric function  $\text{comp}_n(\{a_1..a_m\} - \{b_1..b_k\})$  whose argument is the difference of two hybrid sets each given by ellipsis.

The following expansion can be found in [3].

**Theorem 13** *The complementary symmetric function is given by the following sum*

$$\text{comp}_n(a_1, a_2, \dots | b_1, b_2, \dots) = \sum_{0 < \alpha_1 < \alpha_2 < \dots < \alpha_n} \prod_{i=1}^n (b_{\alpha_i+1-i} - a_{\alpha_i})$$

*over strictly increasing sequences of positive integers.*

*Proof:* Apply Theorem 12 until all complementary symmetric function disappear either by reducing their degree to zero, or by eliminating all their variables.  $\square$

## 2.2 Connection Constants

In order to state the main result of this section, we must first introduce the concept of persistent roots.

**Definition 14** *Let  $(b_n)_{n \in \mathbb{Z}}$  be a sequence of constants indexed by an integer, and let*

$$q_n(x) = \prod_{i=1}^n (x - b_i)$$

*for all  $n$  positive or negative. In other words,*

$$\begin{array}{ccc} \vdots & & \vdots \\ q_2(x) & = & (x - b_1)(x - b_2) \\ q_1(x) & = & (x - b_1) \\ q_0(x) & = & 1 \\ q_{-1}(x) & = & 1/(x - b_0) \\ q_{-2}(x) & = & 1/(x - b_0)(x - b_{-1}) \\ \vdots & & \vdots \end{array}$$

*Then  $(q_n(x))_{n \in \mathbb{Z}}$  is called the sequence of rational functions with persistent roots  $(b_n)_{n \in \mathbb{Z}}$ .*

We can now state the main result concerning the complementary symmetric function.

**Theorem 15** Let  $f(x)$  be a monic rational function of degree  $n$ , and  $(b_k)_{k \leq n}$  a sequence of constants indexed by an integer. Then  $f(x)$  can be expanded in terms of the sequence of rational functions  $(q_k(x))_{k \leq n}$  with persistent roots  $(b_k)_{k \leq n}$  as follows:

$$f(x) = \sum_{k=0}^{\infty} \text{comp}_k(\text{Roots}(f) - \{b_1..b_{n-k+1}\})q_{n-k}(x). \quad (5)$$

Notes to Theorem 15:

1. The degree of a rational function  $f(x) = p(x)/q(x)$  is defined to be the degree of the numerator  $p(x)$  minus the degree of the denominator  $q(x)$ .
2. For  $0 \leq k \leq n$ , the coefficient of  $q_k(x)$  does not depend on  $b_{k+2}, b_{k+3}, \dots, b_n$  even though initial consideration of the problem might seem to indicate that it should.
3. To interpret this theorem when  $k > n$ , recall the definition of improper intervals from section 3.
4. The coefficient of  $q_n(x)$  does not depend on the non-existent " $b_{n+1}$ ," since  $\text{comp}_0(V)$  is identically one.
5. The case in which  $f(x)$  is a polynomial (see Corollary 17) is much simpler and is a direct consequence of equations 1, 2, 3, 4, and Lemma 10.

*Proof of Theorem 15:* We proceed by induction on the number of roots and poles which  $f(x)$  possesses. The base case is thus  $f(x) = 1$ . In this case, the right hand side of equation 5 is

$$\sum_{k=0}^{\infty} \text{comp}_k(-\{b_1..b_{1-k}\})q_{-k}(x).$$

For  $k = 0$ , the summand is one, since the complementary symmetric function of degree zero is identically one and  $q_0(x) = 1$ . The remaining terms are all of the form

$$\frac{e_k(-b_0, -b_{-1}, \dots, -b_{2-k})}{(x - b_0)(x - b_{-1}) \cdots (x - b_{1-k})}$$

which is zero for lack of sufficiently many arguments to the elementary symmetric function. Thus, the sum is one as required.

Next, towards induction assume Eq. 5 holds for  $g(x)$ . Now, consider  $f(x) = (x - a)g(x)$ .

$$\begin{aligned} f(x) &= (x - a) \sum_{k=0}^{\infty} \text{comp}_k(\text{Roots}(g) - \{b_1..b_{n-k}\})q_{n-k}(x) \\ &= \sum_{k=0}^{\infty} \text{comp}_k(\text{Roots}(g) - \{b_1..b_{n-k}\})[(x - b_{n-k}) + (b_{n-k} - a)]q_{n-k-1}(x) \\ &= q_n(x) + \sum_{k=0}^{\infty} [\text{comp}_{k+1}(\text{Roots}(g) - \{b_1..b_{n-k-1}\}) \\ &\quad + (b_{n-k} - a)\text{comp}_k(\text{Roots}(g) - \{b_1..b_{n-k}\})]q_{n-k-1}(x) \\ &= q_n(x) + \sum_{k=0}^{\infty} \text{comp}_k(\text{Roots}(f) - \{b_1..b_{n-k}\})q_{n-k-1}(x) \\ &= \sum_{j=0}^{\infty} \text{comp}_j(\text{Roots}(f) - \{b_1..b_{n-j}\})q_{n-j}(x) \end{aligned}$$

where the fourth equality results from the recursion theorem (12).

The proof for  $f(x) = g(x)/(x - a)$  can be found by reading the above sequence of equalities in reverse order.  $\square$

Theorem 15 gives rise to the following inversion formula.



**Corollary 16** Let  $(a_n)_{n \in \mathbb{Z}}$ ,  $(b_n)_{n \in \mathbb{Z}}$ ,  $(c_n)_{n \in \mathbb{Z}}$ , and  $(d_n)_{n \in \mathbb{Z}}$  be sequences of real numbers. Then

$$c_n = \sum_{k=0}^{\infty} \text{comp}_k(\{a_1..a_n\} - \{b_1..b_{n-k+1}\})d_{n-k}$$

for all  $n$ , if and only if

$$d_n = \sum_{k=0}^{\infty} \text{comp}_k(\{b_1..b_n\} - \{1..a_{n-k+1}\})c_{n-k}$$

for all  $n$ .  $\square$

Theorem 15 is particularly easy to state in the case of polynomials.

**Corollary 17** Given constants  $(a_i)_{i=1}^{\infty}$  and  $(b_i)_{i=1}^{\infty}$ , we have

$$(x - a_1) \cdots (x - a_n) = \sum_{k=0}^n \text{comp}_k(a_1, \dots, a_n | b_1, \dots, b_{k+1})(x - b_1) \cdots (x - b_k). \square$$

As special cases, we have Eqs. 1 and 2. Other special cases of particularly interesting connection constants along with their combinatorial interpretation in the section below and throughout the rest of this paper.

## 2.3 Example of Connection Constants

We begin with an example of the expansion of an inverse formal power series since this lies outside of the usual theory. Let us expand  $1/x$  in terms of the sequence of rational functions  $q_n(x)$  with persistent poles  $b_0, b_{-1}, b_{-2}, \dots$ ; that is, expand  $1/x$  with respect to

$$\begin{aligned} q_{-1}(x) &= 1/(x - b_0), \\ q_{-2}(x) &= 1/(x - b_0)(x - b_{-1}), \\ q_{-3}(x) &= 1/(x - b_0)(x - b_{-1})(x - b_{-2}), \text{ etc.} \end{aligned}$$

By theorem 15, we can write

$$\begin{aligned} \frac{1}{x} &= \sum_{k=0}^{\infty} \text{comp}_k(|0, b_0, b_{-1}, \dots, b_{1-k})q_{-1-k}(x) \\ &= \sum_{k=0}^{\infty} e_k(-b_0, -b_{-1}, \dots, b_{1-k})q_{-1-k}(x) \\ &= \sum_{k=0}^{\infty} (-1)^k b_0 b_{-1} \cdots b_{1-k} q_{-1-k}(x). \end{aligned}$$

So in particular,

$$\frac{1}{x} = \sum_{n=1}^{\infty} \frac{1}{(x+1)^n}$$

for  $x > 0$  as the convergent sum of a geometric series. Similarly,

$$\frac{1}{x} = \sum_{n=1}^{\infty} \frac{(k-1)!}{(x+1)(x+2) \cdots (x+k)}. \quad (6)$$

Using the classical method of telescoping series, equation 6 can be validated for  $x$  a positive integer. However, to demonstrate this identity in general for  $x > 0$ , the preceding theory of connection constants is necessary.

Other examples of connection constants can be found in the following sections.

### 3 Binomial Coefficients

#### 3.1 Connection Constants

Another classical example of connection constants is that of the binomial coefficient  $\binom{n}{k}$  which is given by the coefficients of the polynomial  $(x+1)^k$ , or more generally the Gaussian coefficients  $\binom{n}{k}_q$  which is given by the coefficients of the polynomial  $(x-1)(x-q)(x-q^2)\cdots(x-q^{n-1})$  (see Figure 2 and 3). These constants then have well known combinatorial interpretations.  $\binom{n}{k}$  is the number of  $k$ -element subsets of a given  $n$ -element set whereas the coefficient of  $q^t$  in  $\binom{n}{k}_q$  counts the number of partitions  $\lambda \vdash t$  of length  $k$  with all parts distinct and less than or equal to  $n$  where a partition of  $t$  is defined as a nonincreasing sequence of nonnegative integers summing to  $t$ .<sup>1</sup>

Now, that we are capable of calculating these connection constants via Theorem 15, we can equally well interpret them for  $n$  negative or positive. We are then calculating the coefficient of  $x^k$  in an inverse Laurent series.

By Theorem 12, we have the recurrences

$$\binom{n}{k}_q = \binom{n-1}{k-1}_q + q^{n-1} \binom{n-1}{k}_q \quad (7)$$

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}. \quad (8)$$

Moreover, by theorem 15, we have

$$\begin{aligned} \binom{n}{k}_q &= \text{comp}_{n-k}(-1, -q, \dots, -q^{n-1}) \\ &= \begin{cases} e_{n-k}(1, q, \dots, q_{n-1}) & \text{for } 0 \leq k \leq n, \text{ and} \\ (-1)^{n-k} h_{n-k}(q^{-1}, q^{-2}, \dots, q^n) & \text{for } k \leq n < 0. \end{cases} \\ &= \frac{\prod_{i=1}^k q^{n+1-i} - 1}{(q^k - 1)(q^{k-1} - 1) \cdots (q - 1)}. \end{aligned} \quad (9)$$

And in particular for  $q = 1$ ,

$$\begin{aligned} \binom{n}{k} &= \text{comp}_{n-k}(\underbrace{-1, -1, \dots, -1}_{n \text{ terms}}) \\ &= \begin{cases} e_{n-k}(\underbrace{1, 1, \dots, 1}_{n \text{ terms}}) & \text{for } 0 \leq k \leq n, \text{ and} \\ (-1)^{n-k} h_{n-k}(\underbrace{1, 1, \dots, 1}_{-n \text{ terms}}) & \text{for } k \leq n < 0 \end{cases} \\ &= \frac{n(n-1)(n-2) \cdots (n-k+1)}{k!}. \end{aligned}$$

---

<sup>1</sup>Our definition of the Gaussian coefficient differs from with certain other definitions (for example, see [10]) by a factor of  $q^{(n-k)(n-k-1)/2}$ . Under this alternate definition, the coefficients of  $\binom{n}{k}_q$  then count the number of partitions  $\lambda \vdash t$  of length  $k$  with all parts less than or equal to  $n-k$ . Moreover for  $q$  prime,  $\binom{n}{k}_q$  counts the number of  $k$ -dimensional subvector spaces in an  $n$ -dimension vector spaces over a  $q$ -element field.

Setting all the variables of the elementary or complete symmetric function to one gives the number of terms. In the case of the elementary symmetric function, this is the number of ways of choosing  $k$  variables among  $n$  without replacement. Thus, we have rediscovered the classical combinatorial interpretation of  $\binom{n}{k}$  for  $n$  positive. For the complete symmetric function, the number of terms is the number of ways of choosing  $k$  variables among  $n$  with replacement. In other words, up to sign,  $\binom{-n}{k}$  is the number of  $k$ -element multisets based on a given  $n$ -element set. Hence, we have combinatorial interpretations of  $\binom{n}{k}$  for  $n$  positive or negative.

### 3.2 Roman Coefficients

What is the connection between these two interpretations? And can we permit  $k$  to be negative as well?

We can for example, adopt the following definition of  $\binom{n}{k}$  (consistent with the definition above) for all values of  $n$  and  $k$ , positive or negative (see Figure 1).

$$\binom{n}{k} = \lim_{\epsilon \rightarrow 0} \frac{\Gamma(n+1+\epsilon)}{\Gamma(k+1+\epsilon)\Gamma(n-k+1+\epsilon)}. \quad (10)$$

Table 1: Binomial Coefficients,  $\binom{n}{k}$

| $n \backslash k$ | -4 | -3 | -2 | -1 | 0 | 1  | 2  | 3   | 4  | 5    | 6   |
|------------------|----|----|----|----|---|----|----|-----|----|------|-----|
| 6                | 0  | 0  | 0  | 0  | 1 | 6  | 15 | 20  | 15 | 6    | 1   |
| 5                | 0  | 0  | 0  | 0  | 1 | 5  | 10 | 10  | 5  | 1    | 0   |
| 4                | 0  | 0  | 0  | 0  | 1 | 4  | 6  | 4   | 1  | 0    | 0   |
| 3                | 0  | 0  | 0  | 0  | 1 | 3  | 3  | 1   | 0  | 0    | 0   |
| 2                | 0  | 0  | 0  | 0  | 1 | 2  | 1  | 0   | 0  | 0    | 0   |
| 1                | 0  | 0  | 0  | 0  | 1 | 1  | 0  | 0   | 0  | 0    | 0   |
| 0                | 0  | 0  | 0  | 0  | 1 | 0  | 0  | 0   | 0  | 0    | 0   |
| -1               | -1 | 1  | -1 | 1  | 1 | -1 | 1  | -1  | 1  | -1   | 1   |
| -2               | 3  | -2 | 1  | 0  | 1 | -2 | 3  | -4  | 5  | -6   | 7   |
| -3               | -3 | 1  | 0  | 0  | 1 | -3 | 6  | -10 | 15 | -21  | 28  |
| -4               | 1  | 0  | 0  | 0  | 1 | -4 | 10 | -20 | 35 | -56  | 84  |
| -5               | 0  | 0  | 0  | 0  | 1 | -5 | 15 | -35 | 70 | -126 | 210 |

**Proposition 18 (The Six Regions)** *Let  $n$  and  $k$  be integers. Depending on what region of the Cartesian plane  $(n, k)$  is in, the following formulas apply:*

1. If  $n \geq k \geq 0$ , then  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ .
2. If  $k \geq 0 > n$ , then  $\binom{n}{k} = (-1)^k \binom{-n+k-1}{k}$ .
3. If  $0 > n \geq k$ , then  $\binom{n}{k} = (-1)^{n+k} \binom{-k-1}{n-k}$ .
4. If  $k > n \geq 0$ ,  $\binom{n}{k} = 0$ .
5. If  $0 > k > n$ ,  $\binom{n}{k} = 0$ .
6. If  $k \geq 0 > n$ ,  $\binom{n}{k} = 0$ .  $\square$

Note that in regions 4–6 there is an extra factor of  $\epsilon$  in the numerator of the limit, so we are left with zero. Next, region 1 is the classical case, so we have the usual binomial coefficients. Finally, regions 2 and 3 are simply related to region 1.

Most of the usual properties of binomial coefficients hold true in all six regions.

**Proposition 19 (Complementation)** *For all integers  $n$  and  $m$ ,  $\binom{n}{m} = \binom{n}{n-m}$ .  $\square$*

**Proposition 20 (Iteration)** *For all integers  $i, j$ , and  $k$ ,*

$$\binom{i}{j} \binom{j}{k} = \binom{i}{k} \binom{i-j}{j-k}. \square$$

By equation 8,

**Proposition 21 (Pascal)** *Let  $n$  and  $k$  be integers not both zero, then*

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}. \square$$

Nevertheless,  $\binom{0}{0} = 1$  while  $\binom{-1}{-1} + \binom{-1}{0} = 1 + 1 = 2$ .

The coefficients in Regions 1 and 3 have already been seen to be connection constants; that is to say, they are the coefficients of  $x^{n-k}$  in the inverse power series  $(1+x)^n$ . Those in regions 4–6 are zero. This leaves only region 3, which as we see below, can be interpreted as a set of connection constants in terms of formal power series.

**Proposition 22** *For all integers  $n$ ,  $(x+1)^n$  is given by*

$$\sum_{k=-\infty}^n \binom{n}{k} x^{n-k}$$

*as a formal power series.*

*Proof:* The  $(n-k)$ th derivative of  $(1+x)^n$  evaluated at zero is  $(n-k)! \binom{n}{k}$ .  $\square$

### 3.3 Inclusion of Hybrid Sets

These new binomial coefficients  $\binom{n}{k}$  are always integers, but what do they count? In region 1, they count the number of  $k$ -element subsets of a given  $n$ -element set. We claim that given a suitable generalization of the notion of a subset, this is true in general (up to sign). There is no need for a separate interpretation of  $\binom{n}{k}$  in regions 2 and 3.

We define a partial order on these sets which is a generalization of the usual ordering of classical sets and multisets by inclusion.

**Definition 23 (Subsets)** *Informally, we say  $f$  is a subset of  $g$  (and write  $f \subseteq g$ ) if one can remove elements one at a time from  $g$  (never removing an element that is not a member of  $g$ ) and thus either achieve  $f$  or have removed  $f$ .*

For example, we might start with the hybrid set  $f = \{a, b, c, c|d, e\}$ . We will remove a few of its elements one at a time. Suppose we remove  $b$ ; this leaves  $\{a, c, c|d, e\}$ . Now,  $b$  is no longer an element, so we can not remove it again. Instead, we might remove  $d$  leaving  $\{a, c, c|d, e\}$ . Obviously, we can remove  $d$  as many times as we choose. Finally, suppose we remove  $c$  leaving  $\{a, c|d, d, e\}$ . Hence, we have proven two things. Since we were able to remove  $\{b, d, c\}$ , we know that  $\{b, d, c\} \subseteq f$ . Also, since we were left with  $\{a, c|d, d, e\}$ , we know that  $\{a, c|d, d, e\} \subseteq f$ .

**Proposition 24** [14] *The ordering “ $\subseteq$ ” defined above is a well defined partial order.*  $\square$

*Proof:* See [14].

Note however that the ordering  $\subseteq$  does not form a lattice. For example, hybrid sets  $f = \{|a, b\}$  and  $g = \{a|b\}$  have lower bounds  $\{a|\}$ ,  $\{a, b|\}$ , and  $\{a, b, b|\}$ , but no greatest lower bound. Conversely,  $f$  and  $g$  possess no upper bounds.

**Theorem 25** *The subsets of a classical set  $f$  correspond to the classical subsets of a classical set. The subsets of a multiset  $f$  correspond to the classical submultisets of a multiset.*

*Proof:* To construct a conventional subset of a set  $S$ , we merely remove some elements subject to the conditions that we only remove elements of  $S$  and we don’t remove an element twice. The order of removal is not relevant.

To construct a conventional submultiset of a multiset  $M$ , we merely remove some elements subject to the conditions that we do not remove any element more times than its multiplicity in  $M$ .  $\square$

Now, we have the desired interpretation of the binomial coefficients.

**Theorem 26** *Let  $n$  and  $k$  be arbitrary integers. Let  $f$  be an  $n$ -element new set. Then  $|\binom{n}{k}|$  counts the number of  $k$ -element hybrid sets which are subsets of  $f$ .*  $\square$

Let us consider this result in each of the six regions mentioned in Proposition 18.

1. This is the only classical case. In this region, one might count the number of 2-element subsets of the set  $\{a, b, c, d\}$ . By Theorem 25, we enumerate the usual subsets:  $\{a, b\}$ ,  $\{a, c\}$ ,  $\{a, d\}$ ,  $\{b, c\}$ ,  $\{b, d\}$ , and  $\{c, d\}$  but no others.
2. In this region, one might count the number of 2-element subsets of  $f = \{a, b, c\}$ . These subsets correspond to what we can remove from  $f$ , since what we would have left over after a removal would necessarily contain a negative number of elements. We can remove any of the three elements any number of times, so we have:  $\{a, a\}$ ,  $\{b, b\}$ ,  $\{c, c\}$ ,  $\{a, b\}$ ,  $\{b, c\}$ , and  $\{a, c\}$ .
3. Here, we are interested in  $-5$ -element subsets of  $f$ . Since  $f$  contains  $-3$  elements, we must start with  $f$  and remove 2 elements. Thus, there is one subset here for each subset in the corresponding position in region 2. In this case they are:  $\{a, a, a, b, c\}$ ,  $\{a, b, b, b, c\}$ ,  $\{a, b, c, c, c\}$ ,  $\{a, a, b, b, c\}$ ,  $\{a, b, b, c, c\}$ , and  $\{a, a, b, c, c\}$ .
4. By Theorem 25, there are no 6-element subsets of the set  $\{a, b, c, d\}$ . Once you remove 4 elements, you can not remove any more.
5. There are no  $-2$ -element subset of a  $-3$ -element set  $f$ . If we remove elements from  $f$ , we are left with less than  $-3$  elements, and have removed a positive number of elements. In neither case have we qualified a  $-2$ -element hybrid set to be a subset of  $f$ .
6. Again by Theorem 25, there are no  $-2$ -element subsets of the set  $\{a, b, c, d\}$ , since you are not allowed to introduce elements with negative multiplicities.

## 4 Linear Partitions

Table 2: Gaussian Coefficients,  $\binom{n}{k}_q$ —Region 1

| $n \setminus k$ | 0        | 1                                     | 2                                                 | 3                                               | 4                              |
|-----------------|----------|---------------------------------------|---------------------------------------------------|-------------------------------------------------|--------------------------------|
| 5               | $q^{10}$ | $q^6 + q^7 + q^8$<br>$+ q^9 + q^{10}$ | $q^3 + q^4 + 2q^5 + 2q^6$<br>$+ 2q^7 + q^8 + q^9$ | $q + q^2 + 2q^3 + 2q^4$<br>$+ 2q^5 + q^6 + q^7$ | $1 + q + q^2$<br>$+ q^3 + q^4$ |
| 4               | $q^6$    | $q^3 + q^4 + q^5 + q^6$               | $q + q^2 + 2q^3 + q^4 + q^5$                      | $1 + q + q^2 + q^3$                             | 1                              |
| 3               | $q^3$    | $q + q^2 + q^3$                       | $1 + q + q^2$                                     | 1                                               | 0                              |
| 2               | $q$      | $1 + q$                               | 1                                                 | 0                                               | 0                              |
| 1               | 1        | 1                                     | 0                                                 | 0                                               | 0                              |
| 0               | 1        | 0                                     | 0                                                 | 0                                               | 0                              |

Table 3: Gaussian Coefficients,  $\binom{n}{k}_q$ —Region 2


| $n \setminus k$ | -5                                            | -4                          | -3                 | -2        | -1 |
|-----------------|-----------------------------------------------|-----------------------------|--------------------|-----------|----|
| -1              | $q^{-4}$                                      | $-q^{-3}$                   | $q^{-2}$           | $-q^{-1}$ | 1  |
| -2              | $-q^{-6} - q^{-5} - q^{-4} - q^{-3}$          | $q^{-4} + q^{-3} + q^{-2}$  | $-q^{-2} - q^{-1}$ | 1         | 0  |
| -3              | $q^{-6} + q^{-5} + 2q^{-4} + q^{-3} + q^{-2}$ | $-q^{-3} - q^{-2} - q^{-1}$ | 1                  | 0         | 0  |
| -4              | $-q^{-4} - q^{-3} - q^{-2} - q^{-1}$          | 1                           | 0                  | 0         | 0  |
| -5              | 1                                             | 0                           | 0                  | 0         | 0  |

Just as we did with the binomial coefficients, we must now interpret our generalized Gaussian coefficients according to their combinatorial properties. We will do so in terms of *linear partitions*. One usually defines a (linear) partition or a (linear) partition with distinct parts to be a finite sequence  $\lambda$  of nonnegative integers with  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$  or  $\lambda_1 > \lambda_2 > \dots > \lambda_k$  respectively. Since we are now giving negative integers a status equal to that of nonnegative ones, the classical definition becomes inappropriate. We now proceed to generalize the notion of a partition by first generalizing the relations  $\geq$  and  $>$ .

**Definition 27 (Hybrid Inequalities)** Given integers  $i$  and  $j$ . We say that  $i \ll j$  or equivalently  $j \gg i$  if and only if  $i \in \{0, 1, \dots, j-1\}$ . Similarly, we say that  $i \leq j$  or equivalently  $j \geq i$  if and only if  $j \in \{1, 2, \dots, j\}$ .

Note that these relations have their usual sense for positive integers, but behave rather oddly for negative integers. For example,  $-2, -1, 0$  are the integers  $\leq -3$ , and  $-3, -2, -1$  are the integers  $\ll -3$ . These relations are not partial orders since they do not obey reflexivity. However, they do obey antisymmetry and transitivity, so we can still discuss their Hasse diagrams. As you can see in figure 4, the negative integers

Figure 1: Hasse diagrams of  $\ll$  and  $\leq$



are now unrelated to the positive integers,<sup>2</sup> and the usual order of negative integers has been reversed.

**Definition 28 (New Partitions)** Given a sequence  $\lambda$  of integers, we say that  $\lambda$  is an  $n$ -partition of length  $k$  (and width  $n$ ) if  $(n \gg) \lambda_1 \gg \lambda_2 \gg \dots \gg \lambda_k$ . We say that  $\lambda$  is a  $d$ -partition of length  $k$  (and width  $n$ ) if  $(n \gg) \lambda_1 \gg \lambda_2 \gg \dots \gg \lambda_k$ .

If  $t$  is the sum of the  $k$  parts of  $\lambda$  then we write  $\lambda \vdash t$  or  $|\lambda| = t$ , and we say  $\lambda$  is an  $n$ -partition or  $d$ -partition of  $t$ . The length of  $\lambda$  is denoted  $\ell(\lambda)$ .

By considering the case  $t$  positive, we see that the  $n$ -partition generalizes the idea of a **normal** partition and the  $d$ -partition generalizes the idea of a partition with **distinct** parts.<sup>3</sup>

For example,  $(5, 3, 2, 0)$  is a  $d$ -partition of 10 with length 4 and minimum width 6.  $(5, 5, 3, 2)$  is an  $n$ -partition of 15 with length 4 and minimum width 5.  $(-32, -7, -7)$  is a  $d$ -partition of  $-46$  with length 3 and minimum (maximum?) width  $-32$ .  $(-12, -7, -6, -3, 0)$  is an  $n$ -partition of  $-22$  with length 5 and minimum (maximum?) width  $-13$ . Note also that a sequence can be at the same time a  $d$ -partition and an  $n$ -partition, for example  $(5, 3, 2)$ .

It is well known (see for example [9]) that for  $n$  and  $k$  nonnegative,  $\binom{n}{k}_q$  is a monic polynomial. Its coefficient of  $q^t$  is the number of partitions  $\lambda$  of  $t$  of length  $n - k$  (or equivalently  $k$ ) with all parts less than or equal to  $n$ .

In general, we now have:

**Theorem 29** Let  $k$  be a nonnegative integer, and  $n$  and  $t$  be arbitrary integers. Then  $\binom{n}{k}_q = \sum_t c_t q^t$  where  $c_t$  is the number of  $d$ -partitions  $\lambda$  of  $t$  with length  $k$ , and width  $n$ .

*Proof:* By equation 9.  $\square$

We can now express the  $q$ -binomial coefficients of regions 1 and 3 in terms of each other.

**Corollary 30** Let  $n$  and  $k$  be nonnegative integers, then

$$\binom{n}{k}_q = (-1)^{n+1} q^{\binom{n+1}{2} - \binom{k+1}{2}} \binom{-k-1}{-n-1}_q. \square$$

<sup>2</sup>  $\ll$  and  $\leq$  disagree on whether zero is positive or negative.

<sup>3</sup> Note however that for  $t$  negative, it is the  $n$ -partitions which have distinct parts instead of the  $d$ -partitions.

## 5 Stirling Numbers

### 5.1 Classical Region

Our next example of connection constants is that of Stirling number of the first and second kind,  $s(n, k)$

Table 4: Stirling Numbers of the First Kind,  $s(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1    | 2   | 3    | 4   | 5   | 6 |
|------------------|---|------|-----|------|-----|-----|---|
| 6                | 0 | -120 | 274 | -225 | 85  | -15 | 1 |
| 5                | 0 | 24   | -50 | 35   | -10 | 1   | 0 |
| 4                | 0 | -6   | 11  | -6   | 1   | 0   | 0 |
| 3                | 0 | 2    | -3  | 1    | 0   | 0   | 0 |
| 2                | 0 | -1   | 1   | 0    | 0   | 0   | 0 |
| 1                | 0 | 1    | 0   | 0    | 0   | 0   | 0 |
| 0                | 1 | 0    | 0   | 0    | 0   | 0   | 0 |

and  $S(n, k)$  (see Figures 4 and 5), relating the nonnegative powers of  $x$   $(x^n)_{n \geq 0}$  and the lower factorial

Table 5: Stirling Numbers of the Second Kind,  $S(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1 | 2  | 3  | 4  | 5  | 6 |
|------------------|---|---|----|----|----|----|---|
| 6                | 0 | 1 | 31 | 90 | 65 | 15 | 1 |
| 5                | 0 | 1 | 15 | 25 | 10 | 1  | 0 |
| 4                | 0 | 1 | 7  | 6  | 1  | 0  | 0 |
| 3                | 0 | 1 | 3  | 1  | 0  | 0  | 0 |
| 2                | 0 | 1 | 1  | 0  | 0  | 0  | 0 |
| 1                | 0 | 1 | 0  | 0  | 0  | 0  | 0 |
| 0                | 1 | 0 | 0  | 0  | 0  | 0  | 0 |

polynomials  $((x)_n)_{n \geq 0} = (x(x-1) \cdots (x-n+1))_{n \geq 0}$

$$(x)_n = \sum_{k=0}^n s(n, k) x^k,$$

$$x^n = \sum_{k=0}^n S(n, k) (x)_k.$$

Their best known combinatorial interpretations is that  $S(n, k)$  counts the number of partitions of an  $n$ -element set into  $k$  blocks while (up to sign)  $s(n, k)$  counts the number of partitions of an  $n$ -element set containing  $k$ -cycles.

When generalizing these constants to negative values of  $n$ , we define the lower factorial sequence of rational functions  $((x)_n)_{n \in \mathbb{Z}}$  for  $n$  an arbitrary integer as the sequence of rational functions with persistent roots  $(i-1)_{i \in \mathbb{Z}}$ . Next, we must decide as we did in Proposition 22 whether to consider expansion of  $(x)_n$  as a formal power series or as an inverse Laurent series. In Proposition 22, the difference was not critical; the binomial coefficients in regions 2 and 3 are related by Proposition 19. Each region is the reflection of the other.



Table 6: Stirling Numbers of the First Kind  $s(n, k)$ —Region 2

| $k \backslash n$ | -5                                          | -4                              | -3                        | -2                | -1 |
|------------------|---------------------------------------------|---------------------------------|---------------------------|-------------------|----|
| 6                | $\frac{226,576,031,859}{5,598,720,000,000}$ | $\frac{11,679,655}{71,663,616}$ | $\frac{137,845}{279,936}$ | $\frac{127}{128}$ | 1  |
| 5                | $-\frac{3,673,451,957}{93,312,000,000}$     | $-\frac{952,525}{5,971,968}$    | $-\frac{22,631}{46,656}$  | $-\frac{63}{64}$  | -1 |
| 4                | $\frac{58,067,611}{1,555,200,000}$          | $\frac{76,111}{497,664}$        | $\frac{3661}{7776}$       | $\frac{31}{32}$   | 1  |
| 3                | $-\frac{874,853}{25,920,000}$               | $-\frac{5845}{41,472}$          | $-\frac{575}{1296}$       | $-\frac{15}{16}$  | -1 |
| 2                | $\frac{12,019}{432,000}$                    | $\frac{415}{3456}$              | $\frac{85}{216}$          | $\frac{7}{8}$     | 1  |
| 1                | $-\frac{137}{7200}$                         | $-\frac{25}{288}$               | $-\frac{11}{36}$          | $-\frac{3}{4}$    | -1 |
| 0                | $\frac{1}{120}$                             | $\frac{1}{24}$                  | $\frac{1}{6}$             | $\frac{1}{2}$     | 1  |

## 5.2 Formal Power Series Region

Here, as we will see, there is no such symmetry. First, let us examine regions 1 and 2 defined by formal power series (see table 6). These constants were considered in [15, 13]. However, except for the classical region 1, we no longer have integers, so there is no combinatorial interpretation to be expected. However, we do have several interesting recursions and identities which prove to be useful in [11] such as

**Theorem 31** [13, Porism 4.2 and Proposition 4.12] *For all integers  $n$  (positive or negative) and nonnegative integers  $k$ ,*

$$s(n, k) = \lim_{\epsilon \rightarrow 0} \Gamma(1 - n + \epsilon)^{-1} \text{comp}_k \left( \frac{1}{\epsilon}, \frac{1}{-1 + \epsilon}, \frac{1}{-2 + \epsilon}, \dots, \frac{1}{-n + \epsilon} \right). \square$$

We also have the recursion

**Theorem 32** [13, Theorem 3.2] *For all integers  $n$  and for all positive integers  $k$ ,*

$$s(n + 1, k) = s(n, k - 1) - ns(n, k). \square$$

In region 2, we have the following result due to Knuth.

**Proposition 33** [13, Proposition 4.3] *Let  $n$  and  $k$  be nonnegative integers (not both zero). Then  $s(n, k)$  is given by the following finite sum:*

$$s(-n, k) = \frac{(-1)^{k+1}}{n!} \sum_{m=1}^n \binom{n}{m} (-1)^m m^{-k}. \quad (11)$$

Note that Proposition 33 is the analog of the following classical result [2] involving Stirling numbers of the *second* kind  $S(k, n)$ .

$$S(k, n) = \frac{(-1)^n}{n!} \sum_{m=1}^k \binom{n}{m} (-1)^m m^k.$$

Thus, in some sense to be made clear later, we can say that

$$S(k, n) = (-1)^{n+k+1} s(-n, -k).$$

Table 7: Stirling Numbers of the First Kind,  $s(n, k)$ —Region 3

| $n \backslash k$ | -6  | -5  | -4 | -3 | -2 | -1 |
|------------------|-----|-----|----|----|----|----|
| -1               | -1  | 1   | -1 | 1  | -1 | 1  |
| -2               | 31  | -15 | 7  | -3 | 1  | 0  |
| -3               | -90 | 25  | -6 | 1  | 0  | 0  |
| -4               | 65  | -10 | 1  | 0  | 0  | 0  |
| -5               | -15 | 1   | 0  | 0  | 0  | 0  |
| -6               | 1   | 0   | 0  | 0  | 0  | 0  |

Table 8: Stirling Numbers of the Second Kind,  $S(n, k)$ —Region 3

| $n \backslash k$ | -6  | -5 | -4 | -3 | -2 | -1 |
|------------------|-----|----|----|----|----|----|
| -1               | 120 | 24 | 6  | 2  | 1  | 1  |
| -2               | 274 | 50 | 11 | 3  | 1  | 0  |
| -3               | 225 | 35 | 6  | 1  | 0  | 0  |
| -4               | 85  | 10 | 1  | 0  | 0  | 0  |
| -5               | 15  | 1  | 0  | 0  | 0  | 0  |
| -6               | 1   | 0  | 0  | 0  | 0  | 0  |

### 5.3 Inverse Laurent Series Region

On the other hand, in the case of inverse Laurent series, we can apply the results of section 2 to calculate  $s(n, k)$  and  $S(n, k)$ . See tables 7 and 8.

$$\begin{aligned}
s(n, k) &= \text{comp}_{n-k}(0..n-1) \\
&= \begin{cases} (-1)^{n-k} e_{n-k}(1, 2, \dots, n-1) & \text{for } n \geq k \geq 0, \\ h_{n-k}(-1, -2, \dots, n) & \text{for } 0 > n \geq k. \end{cases} \\
S(n, k) &= \text{comp}_{n-k}(-\{0..k\}) \\
&= \begin{cases} h_{n-k}(1, 2, \dots, k) & \text{for } n \geq k \geq 0, \\ (-1)^k e_{n-k}(-1, -2, \dots, k+1) & \text{for } 0 > n \geq k. \end{cases}
\end{aligned}$$

By Theorem 12, the numbers in regions 1 and 3 obey the identities

$$\begin{aligned}
s(n, k) &= s(n-1, k-1) - ns(n-1, k) \\
S(n, k) &= S(n-1, k-1) + kS(n-1, k).
\end{aligned}$$

These Stirling numbers are always integers, so we can hope to give an extended combinatorial interpretation. We will not only do that, but in fact, we will give a combinatorial interpretation of the extended  $p, q$ -Stirling numbers. The  $p, q$ -Stirling numbers of the first kind (see tables 9 and 10) and second kind  $s_{pq}(n, k)$  and  $S_{pq}(n, k)$  (see tables 11 and 12) are the coefficients connecting the powers of  $x$  with the  $p, q$ -lower factorial  $(x; p, q)_n$  where the  $p, q$ -lower factorials  $((x; p, q)_n)_{n \in \mathbb{Z}}$  form the sequence of rational functions with persistent roots given by the  $p, q$ -numbers  $(\sqrt{pq}(p^i - q^i)/(p - q))_{i \in \mathbb{Z}}$ .<sup>4</sup>

<sup>4</sup>The  $p, q$ -numbers are sometimes defined as  $(p^i - q^i)/(p - q)$  or simply  $p^i - q^i$ . These alternate definitions lead to quite similar results.

Table 9:  $p, q$ -Stirling Numbers of the First Kind,  $s_{pq}(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1                                          | 2                                                              | 3                         | 4 |
|------------------|---|--------------------------------------------|----------------------------------------------------------------|---------------------------|---|
| 4                | 0 | $-\sqrt{pq}^3 [q^3 + 2q^2p + 2qp^2 - p^3]$ | $pq [ (q + q^2 + q^3) + (1 + q + 2q^2)p + (1 + 2q)p^2 + p^3 ]$ | $-\sqrt{pq}(1 + q + q^2)$ | 1 |
| 3                | 0 | $pq[q + p]$                                | $-\sqrt{pq}(1 + p + q)$                                        | 1                         | 0 |
| 2                | 0 | $-\sqrt{pq}$                               | 1                                                              | 0                         | 0 |
| 1                | 0 | 1                                          | 0                                                              | 0                         | 0 |
| 0                | 1 | 0                                          | 0                                                              | 0                         | 0 |

 Table 10:  $p, q$ -Stirling Numbers of the First Kind,  $s_{pq}(n, k)$ —Region 3

| $n \backslash k$ | -4                                                                 | -3                                   | -2             | -1 |
|------------------|--------------------------------------------------------------------|--------------------------------------|----------------|----|
| -1               | $-1/\sqrt{pq}^3$                                                   | $1/pq$                               | $-1/\sqrt{pq}$ | 1  |
| -2               | $[p^{-2} + (q^{-1} + 2)p^{-1} + (1 + q^{-1} + q^{-2})]/pq$         | $-[p^{-1} + (q^{-1} + 1)]/\sqrt{pq}$ | 1              | 0  |
| -3               | $-[p^{-2} + (q^{-1} + 1)p^{-1} + (q^{-2} + q^{-1} + 1)]/\sqrt{pq}$ | 1                                    | 0              | 0  |
| -4               | 1                                                                  | 0                                    | 0              | 0  |

 Table 11:  $p, q$ -Stirling Numbers of the Second Kind,  $S_{pq}(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1             | 2                                        | 3                                              | 4 |
|------------------|---|---------------|------------------------------------------|------------------------------------------------|---|
| 4                | 0 | $\sqrt{pq}^3$ | $pq [ (1 + q + q^2) + (1 + 2q)p + p^2 ]$ | $\sqrt{pq} [ (1 + q + q^2) + (1 + q)p + p^2 ]$ | 1 |
| 3                | 0 | $pq$          | $\sqrt{pq}(1 + q + p)$                   | 1                                              | 0 |
| 2                | 0 | $\sqrt{pq}$   | 1                                        | 0                                              | 0 |
| 1                | 0 | 1             | 0                                        | 0                                              | 0 |
| 0                | 1 | 0             | 0                                        | 0                                              | 0 |

 Table 12:  $p, q$ -Stirling Numbers of the Second Kind,  $S_{pq}(n, k)$ —Region 3

| $n \backslash k$ | -4                                                                                              | -3                                  | -2            | -1 |
|------------------|-------------------------------------------------------------------------------------------------|-------------------------------------|---------------|----|
| -1               | $[p^{-3} + 2q^{-1}p^{-2} + 2q^{-2}p^{-1} + q^{-3}]/\sqrt{pq}^3$                                 | $[p^{-1} + q^{-1}]/pq$              | $1/\sqrt{pq}$ | 1  |
| -2               | $[p^{-3} + (2q^{-1} + 1)p^{-2} + (2q^{-2} + q^{-1} + 1)p^{-1} + (q^{-3} + q^{-2} + q^{-1})]/pq$ | $[p^{-1} + (q^{-1} + 1)]/\sqrt{pq}$ | 1             | 0  |
| -3               | $[p^{-2} + (q^{-1} + 1)p^{-1} + (q^{-2} + q^{-1} + 1)]/\sqrt{pq}$                               | 1                                   | 0             | 0  |
| -4               | 1                                                                                               | 0                                   | 0             | 0  |

The  $p, q$ -Stirling number obey the following properties:

$$\begin{aligned} s_{pq}(n, k) &= \text{comp}_{n-k}([0]_{pq} \dots [n-1]_{pq}) \\ &= \begin{cases} (-1)^{n-k} e_{n-k}([1]_{pq}, [2]_{pq}, \dots, [n-1]_{pq}) & \text{for } n \geq k \geq 0, \\ h_{n-k}([-1]_{pq}, [-2]_{pq}, \dots, [n]_{pq}) & \text{for } 0 > n \geq k. \end{cases} \end{aligned} \quad (12)$$

$$\begin{aligned} S_{pq}(n, k) &= \text{comp}_{n-k}(-\{[0]_{pq} \dots [k]_{pq}\}) \\ &= \begin{cases} h_{n-k}([1]_{pq}, [2]_{pq}, \dots, [k]_{pq}) & \text{for } n \geq k \geq 0, \\ (-1)^k e_{n-k}([-1]_{pq}, [-2]_{pq}, \dots, [k+1]_{pq}) & \text{for } 0 > n \geq k. \end{cases} \end{aligned} \quad (13)$$

$$\begin{aligned} s_{pq}(n, k) &= s_{pq}(n-1, k-1) - [n]_{pq} s_{pq}(n-1, k) \\ S_{pq}(n, k) &= S_{pq}(n-1, k-1) + [k]_{pq} S_{pq}(n-1, k). \end{aligned}$$

By definition, the  $p, q$ -Stirling numbers transform into the classical Stirling numbers by setting  $p$  and  $q$  equal to one. If we merely set  $p$  equal to one, we have the usual  $q$ -Stirling numbers  $s_q(n, k)$  (see tables

Table 13:  $q$ -Stirling Numbers of the First Kind,  $s_q(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1                                            | 2                                               | 3                                  | 4                       |
|------------------|---|----------------------------------------------|-------------------------------------------------|------------------------------------|-------------------------|
| 5                | 0 | $q^2 + 3q^3 + 5q^4 + 6q^5$                   | $-[4q^{3/2} + 9q^{5/2} + 12q^{7/2} + 12q^{9/2}$ | $6q + 9q^2 + 9q^3$                 | $-[4q^{1/2} - 3q^{3/2}$ |
| 4                | 0 | $+5q^6 + 3q^7 + q^8$                         | $8q^{11/2} + 4q^{13/2} + q^{15/2}]$             | $+7q^4 + 3q^5 + q^6$               | $+2q^{5/2} + q^{7/2}]$  |
| 3                | 0 | $-[q^{3/2} + 2q^{5/2} + 2q^{7/2} + q^{9/2}]$ | $3q + 4q^2 + 3q^3 + q^4$                        | $-[3q^{1/2} + 2q^{3/2} + q^{5/2}]$ | 1                       |
| 2                | 0 | $q + q^2$                                    | $-[2q^{1/2} + q^{3/2}]$                         | 1                                  | 0                       |
| 1                | 0 | $-q^{1/2}$                                   | 1                                               | 0                                  | 0                       |
| 0                | 1 | 0                                            | 0                                               | 0                                  | 0                       |

Table 14:  $q$ -Stirling Numbers of the First Kind,  $s_q(n, k)$ —Region 3

| $n \backslash k$ | -5                           | -4                        | -3            | -2               | -1 |
|------------------|------------------------------|---------------------------|---------------|------------------|----|
| -1               | $q^{-2}$                     | $-\sqrt{q}^{-5}$          | $q^{-1}$      | $-\sqrt{q}^{-1}$ | 1  |
| -2               | $-[q^{-9/2} + 4q^{-7/2}$     | $q^{-3} + 3q^{-2}$        | $-[q^{-3/2}$  | 1                | 0  |
| -3               | $+6q^{-5/2} + 4q^{-3/2}]$    | $+3q^{-1}$                | $-2q^{-1/2}]$ | 0                | 0  |
| -4               | $q^{-5} + 3q^{-4} + 7q^{-3}$ | $-[4q^{-5/2} + 2q^{-3/2}$ | 1             | 0                | 0  |
| -5               | $+8q^{-2} + 6q^{-1}$         | $+3q^{-1/2}]$             | 0             | 0                | 0  |
| -6               | $-[q^{-7/2} - 2q^{-5/2}$     | 1                         | 0             | 0                | 0  |
| -7               | $-3q^{-3/2} - 4q^{-1/2}$     | 0                         | 0             | 0                | 0  |

13 and 14) and  $S_q(n, k)$  (see tables 15 and 16) defined as the coefficients connecting the powers of  $x$  with the  $q$ -lower factorial  $(x; q)_n$  where the  $q$ -lower factorials  $((x; q)_n)_{n \in \mathbb{Z}}$  form the sequence of rational functions with persistent roots given by the  $q$ -numbers  $\sqrt{q}(q^i - 1)/(q - 1)$ .

Note also that the  $p, q$ -numbers are invariant under the interchange of  $p$  with  $q$ . Thus, the  $p, q$ -Stirling numbers are symmetric in  $p$  and  $q$ .

This interpretation will be made in terms of Ferrers diagrams.

Table 15:  $q$ -Stirling Numbers of the Second Kind,  $S_q(n, k)$ —Region 1

| $n \backslash k$ | 0 | 1         | 2                                              | 3                                   | 4                                              |
|------------------|---|-----------|------------------------------------------------|-------------------------------------|------------------------------------------------|
| 5                | 0 | $q^2$     | $4q^{3/2} + 6q^{5/2}$<br>$+4q^{7/2} + q^{9/2}$ | $6q + 8q^2 + 7q^3$<br>$+3q^4 + q^5$ | $4q^{1/2} + 3q^{3/2}$<br>$+2q^{5/2} + q^{7/2}$ |
| 4                | 0 | $q^{3/2}$ | $3q + 3q^2 + q^3$                              | $3q^{1/2} + 2q^{3/2} + q^{5/2}$     | 1                                              |
| 3                | 0 | $q$       | $2q^{1/2} + q^{3/2}$                           | 1                                   | 0                                              |
| 2                | 0 | $q^{1/2}$ | 1                                              | 0                                   | 0                                              |
| 1                | 0 | 1         | 0                                              | 0                                   | 0                                              |
| 0                | 1 | 0         | 0                                              | 0                                   | 0                                              |

Table 16:  $q$ -Stirling Numbers of the Second Kind,  $S_q(n, k)$ —Region 3

| $n \backslash k$ | -5                                                                                          | -4                                                | -3                        | -2         | -1 |
|------------------|---------------------------------------------------------------------------------------------|---------------------------------------------------|---------------------------|------------|----|
| -1               | $q^{-8} + 3q^{-7} + 5q^{-6} + 6q^{-5}$<br>$+5q^{-4} + 3q^{-3} + q^{-2}$                     | $q^{-9/2} + 2q^{-7/2}$<br>$+2q^{-5/2} + q^{-3/2}$ | $q^{-2} + q^{-1}$         | $q^{-1/2}$ | 1  |
| -2               | $q^{-15/2} + 4q^{-13/2} + 8q^{-11/2} + 12q^{-9/2}$<br>$+12q^{-7/2} + 9q^{-5/2} + 4q^{-3/2}$ | $q^{-4} + 3q^{-3}$<br>$+4q^{-2} + 3q^{-1}$        | $q^{-3/2}$<br>$+q^{-1/2}$ | 1          | 0  |
| -3               | $q^{-6} + 3q^{-5} + 7q^{-4}$<br>$+9q^{-3} + 9q^{-2} + 6q^{-1}$                              | $q^{-5/2} + 2q^{-3/2}$<br>$+3q^{-1/2}$            | 1                         | 0          | 0  |
| -4               | $q^{-7/2} + 2q^{-5/2} + 3q^{-3/2} + 4q^{-1/2}$                                              | 1                                                 | 0                         | 0          | 0  |
| -5               | 1                                                                                           | 0                                                 | 0                         | 0          | 0  |

**Definition 34 (Ferrers Diagram)** The Ferrers diagram of a  $n$ -partition or  $d$ -partition  $\lambda$  with length  $k$  is defined as the following hybrid set of squares  $(i, j)$ :

$$\mathcal{H}(\lambda) = \sum_{i=1}^k \{(1, 1) \dots (1, \lambda_i)\}.$$

The cardinality of  $\mathcal{H}(\lambda)$  is  $|\lambda|$ .

In the case of classical partitions, the above corresponds to the traditional concept of a Ferrers diagram. For example, the Ferrers diagram of  $(3, 2, 0)$  is the positive set  $\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2)\}$ . On the other hand, the Ferrers diagram of  $(-3, -2)$  is the negative set  $\{(1, 0), (1, -1), (1, -2), (2, 0), (2, -1)\}$ .

**Definition 35 (0-1 Tableaux)** A 0-1  $p$ -tableau (resp. a 0-1  $n$ -tableau) is a pair  $\alpha = (\lambda, f)$  where  $\lambda$  is a  $p$ -partition (resp.  $n$ -partition), and  $f$  is a filling of that diagram. That is to say it is a collection of constants equal to 0 or 1 and denoted  $f_{ij}$  for each  $(i, j) \in \mathcal{H}(\lambda)$  subject to the condition that there is exactly one 1 in each row. ie: for all  $1 \leq i \leq \ell(\lambda)$ , there is a unique  $j$  such that  $f_{ij} = 1$ .

The inversion number  $inv(\alpha)$  of a 0-1 tableau  $\alpha = (\lambda, f)$  is one half plus the number of zeros to the left of a one in the diagram, ie: the number of triples  $(i, j_1, j_2)$  such that  $j_1 \ll j_2$ ,  $f_{ij_1} = 0$ , and  $f_{ij_2} = 1$  while taking into account possible negative multiplicities. Similarly, the non-inversion number  $nin(\alpha)$  of a 0-1 tableau is one half plus the number of zeros to the right of a one in the diagram.

We then have the following generalization of [17, eq. (1.4)]:

**Theorem 36** *Let  $n$  and  $k$  be integers. Then the Stirling numbers of the second kind of degree  $n$  and order  $k$  are given by sums*

$$\begin{aligned} s_{pq}(n, k) &= \sum_{\alpha} q^{\text{inv}(\alpha)} p^{n\text{in}(\alpha)} \\ s_q(n, k) &= \sum_{\alpha} q^{\text{inv}(\alpha)} \end{aligned}$$

*over 0-1 tableaux  $\alpha$  of  $d$ -partitions of width  $n$  and length  $n - k$ . Thus,  $s(n, k)$  is the number of such tableaux.*

*The Stirling numbers of the first kind of degree  $n$  and order  $k$  are given by similar sums*

$$\begin{aligned} S_{pq}(n, k) &= \sum_{\alpha} q^{\text{inv}(\alpha)} p^{n\text{in}(\alpha)} \\ S_q(n, k) &= \sum_{\alpha} q^{\text{inv}(\alpha)} p^{n\text{in}(\alpha)} \end{aligned}$$

*this time over 0-1 tableaux  $\alpha$  of  $n$ -partitions of width  $k$ , and length  $n - k$ . Thus,  $S(n, k)$  is the number of such tableaux.*

*Proof:* Equation 12.  $\square$

**Corollary 37** *Let  $n$  and  $k$  be integers. Then we have the following relations*

$$\begin{aligned} S_{pq}(n, k) &= (-1)^{n+k} s_{p^{-1}q^{-1}}(-k, -n) \\ S_q(n, k) &= (-1)^{n+k} s_{q^{-1}}(-k, -n) \\ S(n, k) &= (-1)^{n+k} s(-k, -n). \square \end{aligned}$$

## References

- [1] P. L. BUTZER, M. HAUSS AND M. SCHMIDT, *Factorial Functions and Stirling Numbers of Fractional Orders*, Results in Mathematics, **16**, 16–48 (1989).
- [2] S. CHARALAMBIDES AND J. SINGH, *A Review of Stirling Numbers: Their Generalization and Statistical Application*, Communications in Statistical Theory: Theory and Methods **17(A)** 2533-95 (1988).
- [3] L. COMTET, “Advanced Combinatorics,” Reidel, 1974.
- [4] E. DAMIANI, O. D’ANTONA, AND D. LOEB, *The Complementary Symmetric Function: Connection Constants Using Negative Sets*, submitted to European Journal of Combinatorics.
- [5] O. D’ANTONA AND E. DAMIANI, *Connection Constants*, Rapport Interno no. 54/89, Universita di Milano, Dipartimento di Scienze dell’Informazione.
- [6] H. W. GOULD, *Note on Recurrence Relations for Stirling Numbers*, Publications de l’Institut Mathématique, Nouvelle série, tome 6 (20) 1966, 115-119.
- [7] H. W. GOULD, *The  $q$ -Stirling Numbers of the First and Second Kinds*, Duke Math. J. , **28** (1961) 281–289.
- [8] J. G. HAGEN, *Synopsis der hoheren Mathematik*, Vol. 1, Berlin, 1891.
- [9] D. KNUTH, *Subspaces, Subsets, and Partitions*, Journal of Combinatorial Theory **10** (1971) 178–180.

- [10] P. LEROUX, *Reduced Matrices and  $q$ -Log-Concavity Properties of  $q$ -Stirling Numbers*, Journal of Combinatorial Theory **A**, **54** (1991) 64–84.
- [11] D. LOEB, *The Iterated Logarithmic Algebra*, Advances in Mathematics, **86** (1991) 155–234.
- [12] D. LOEB, *A Generalization of the Binomial Coefficients*, SIAM Journal of Discrete Mathematics, To appear.
- [13] D. LOEB, *A Generalization of the Stirling Numbers*, SIAM Journal of Discrete Mathematics, To appear.
- [14] D. LOEB, *Sets with A Negative Number of Elements*, Advances in Mathematics, **91** (1992) 64–74.
- [15] D. LOEB AND G.-C. ROTA, *Formal Power Series of Logarithmic Type*, Advances in Mathematics, **75** (1989), 1–118.
- [16] I. G. MACDONALD, “Symmetric Functions and Hall Polynomials,” *Oxford Mathematical Monographs*, Clarendon Press, Oxford, 1979.
- [17] A. MÉDICIS AND P. LEROUX, *A Unified Combinatorial Approach for  $q$ - (and  $p, q$ -)Stirling Numbers*, Rapport de recherche no. 126, Département de mathématiques et d’informatique, Université du Québec à Montréal.
- [18] F. REGONATI, *Sui numeri di Whitney di certi reticoli supersolibi finiti*.
- [19] S. ROMAN, *A Generalization of the Binomial Coefficients*, To Appear.